

1. (C) For $4^x 5^y 6^z = 2^{2x+z} 3^z 5^y$ to be a perfect square, the exponent on each prime must be even. That is, y and z must be even. Only choice (C) satisfies this condition.

2. (B) Checking the first few values we find

$$\begin{aligned} u_0 &= 4 \\ u_1 &= f(4) = 5 \\ u_2 &= f(5) = 2 \\ u_3 &= f(2) = 1 \\ u_4 &= f(1) = 4 \\ u_5 &= f(4) = 5. \end{aligned}$$

In general, we see that $u_{4k+j} = u_j$, where k is any integer greater than or equal to zero. Hence, $u_{2002} = u_{4 \cdot 500 + 2} = u_2 = 2$.

3. (B) Let a, b, c be the dimensions of the box, $a \leq b \leq c$. Since $abc = 2002 = 2 \cdot 7 \cdot 11 \cdot 13$, the only possible triples (a, b, c) are $(1, 1, 2002)$, $(1, 2, 1001)$, $(1, 7, 286)$, $(1, 11, 182)$, $(1, 13, 154)$, $(1, 14, 143)$, $(1, 22, 91)$, $(1, 26, 77)$, $(2, 7, 143)$, $(2, 11, 91)$, $(2, 13, 77)$, $(7, 11, 26)$, $(7, 13, 22)$, and $(11, 13, 14)$. Among these, the last triple gives the minimum sum, 38.

4. (E) Multiplying both sides of the given equation by $b(b + 10a)$ yields

$$2ab + 10a^2 + 10b^2 = 2b^2 + 20ab$$

which is equivalent to $(a - b)(5a - 4b) = 0$. Since $a \neq b$, we have $5a = 4b$, or $\frac{a}{b} = 0.8$.

5. (C) There are 16 factors of $2002 = 2 \cdot 7 \cdot 11 \cdot 13$. We examine them and find that only three of them, namely 2, 7, and 14, are 2 less than a perfect square.
6. (B) If m and w are the current numbers of men and women, respectively, then we have

$$\frac{m}{1.05} + \frac{w}{1.20} = \frac{m+w}{1.10}$$

or

$$\frac{m}{w} \cdot \left(\frac{1}{1.05} - \frac{1}{1.10} \right) = \frac{1}{1.10} - \frac{1}{1.20}.$$

It follows that $\frac{m}{w} = \frac{7}{4}$ so that $\frac{m+w}{w} = \frac{11}{4}$, and $\frac{w}{m+w} = \frac{4}{11}$.

7. (A) Let U be the set of all three-digit numbers, let S be the set of three-digit numbers that contain no 2s, and let T be the set of three digit numbers that contain no 3s. Then $S \cap T$ is the set of three-digit numbers containing neither a 2 nor a 3 and $U - (S \cup T)$ is the set of three-digit numbers containing at least one 2 and at least one 3. We have $|U| = 900$, $|S| = |T| = 8 \cdot 9^2$, and $|S \cap T| = 7 \cdot 8^2$. Therefore $|S \cup T| = |S| + |T| - |S \cap T| = 848$, and $|U - (S \cup T)| = 52$.

8. (D) From similar right triangles ACE and ECB , $\frac{EC}{AC} = \frac{CB}{EC}$. Therefore $EC^2 = (AC)(CB) = (1)(25)$ and $EC = 5$. Similarly, $FD^2 = (AD)(DB) = (8)(18)$ and $FD = 12$. If we choose G on FD so that EG is parallel to AB then triangle EGF is right with $EG = GF = 7$. Therefore the hypotenuse $EF = 7\sqrt{2}$.
9. (D) Let the fly be x meters from the ceiling. Then the fly and point P determine a major diagonal of the rectangular parallelepiped having dimensions 1, 8, and x . Therefore, $1^2 + 8^2 + x^2 = 9^2$, and it follows that $x = 4$.
10. (E) The expression is an identity. Note first that

$$\begin{aligned} f_6(x) &= (\sin^2 x)^3 + (\cos^2 x)^3 \\ &= (\sin^2 x + \cos^2 x)(f_4(x) - \sin^2 x \cos^2 x) \\ &= f_4(x) - \sin^2 x \cos^2 x. \end{aligned}$$

Therefore,

$$6f_4(x) - 4f_6(x) = 2f_4(x) + 4\sin^2 x \cos^2 x = 2f_2(x).$$

11. (C)

$$\begin{aligned} &\frac{1}{t_1} + \frac{1}{t_2} + \frac{1}{t_3} + \dots + \frac{1}{t_{2002}} = \\ &\frac{2}{1 \cdot 2} + \frac{2}{2 \cdot 3} + \frac{2}{3 \cdot 4} + \dots + \frac{2}{2002 \cdot 2003} = \\ &\left(\frac{2}{1} - \frac{2}{2}\right) + \left(\frac{2}{2} - \frac{2}{3}\right) + \left(\frac{2}{3} - \frac{2}{4}\right) + \dots + \left(\frac{2}{2002} - \frac{2}{2003}\right) = \\ &\frac{2}{1} - \frac{2}{2003} = \frac{4004}{2003}. \end{aligned}$$

12. (C) Because $n^3 - 8n^2 + 20n - 13 = (n-1)(n^2 - 7n + 13)$, for the value to be prime one factor must equal 1 and the other factor must be prime. For $n-1 = 1$ we must have $n = 2$, and in this case the other factor is the prime 3. So $n = 2$ is a solution. For $n^2 - 7n + 13 = 1$, we have $n^2 - 7n + 12 = 0 = (n-4)(n-3)$, so we must have $n = 3$ or 4, and in each case the other factor is prime (2 and 3, respectively). Therefore $n^3 - 8n^2 + 20n - 13$ is a prime for three positive integer values of n .
13. (D) Since $1^2 + 2^2 + 3^2 + \dots + 18^2 > 2002$, we know that $n \leq 17$. Then note that $1^2 + 2^2 + 3^2 + \dots + 19^2 - 18^2 - 12^2 = 2002$, hence $n = 17$.

14. (D) The real part of the sum is

$$\begin{aligned} -2 + 4 - 6 + \dots + 2000 - 2002 &= -2 + (4 - 6) + \dots + (2000 - 2002) \\ &= -2 \cdot 501 = -1002, \end{aligned}$$

and the imaginary part of the sum is

$$\begin{aligned} 1 - 3 + 5 - \dots - 1999 + 2001 &= 1 + (-3 + 5) + \dots + (-1999 + 2001) \\ &= 1 + 2 \cdot 500 = 1001. \end{aligned}$$

Hence the answer is (D), $-1002 + 1001i$.

15. (C) There are $\binom{2002}{2} = \frac{2002 \cdot 2001}{2 \cdot 1} = 1001 \cdot 2001$ possible pairs that can be drawn. There are 1001^2 pairs of different colored marbles, so $P_d = \frac{1001^2}{1001 \cdot 2001}$. Therefore, $P_s = 1 - P_d = \frac{1000}{2001}$, and $|P_s - P_d| = \frac{1}{2001}$.

16. (C) If the sides of a triangle corresponding to the altitudes 12, 15, and 20 are a , b , and c respectively, then we have

$$12a = 15b = 20c$$

or, dividing by 60,

$$\frac{a}{5} = \frac{b}{4} = \frac{c}{3}.$$

It follows that $a = 5r$, $b = 4r$, and $c = 3r$ and so the triangle is a 3-4-5 right triangle whose largest angle is 90° .

17. (E) Since $\cos^2 x = 1 - \sin^2 x$ and $\sin^2 x = 1 - \cos^2 x$, we have

$$\begin{aligned} \sqrt{\sin^4 x + 4 \cos^2 x} - \sqrt{\cos^4 x + 4 \sin^2 x} &= \sqrt{\sin^4 x - 4 \sin^2 x + 4} \\ &\quad - \sqrt{\cos^4 x - 4 \cos^2 x + 4} \\ &= \sqrt{(\sin^2 x - 2)^2} - \sqrt{(\cos^2 x - 2)^2} \\ &= |\sin^2 x - 2| - |\cos^2 x - 2| \\ &= 2 - \sin^2 x - (2 - \cos^2 x) \\ &= \cos^2 x - \sin^2 x = \cos 2x. \end{aligned}$$

18. (A) Adding the three equations we obtain

$$(a^2 + 6a) + (b^2 + 2b) + (c^2 + 4c) = -14,$$

which is equivalent to

$$(a + 3)^2 + (b + 1)^2 + (c + 2)^2 = 0.$$

Therefore $a = -3$, $b = -1$, $c = -2$, and $a^2 + b^2 + c^2 = 14$.

19. (D) Let the extensions of AB beyond B and DC beyond C meet at E . Then, because angles CBE and BCE both equal 60° , BEC is an equilateral triangle of side 4. The area of triangle BCE is $4\sqrt{3}$ and the area of triangle AED is $\frac{(AE)(DE)\sin 60^\circ}{2} = \frac{63\sqrt{3}}{4}$. Therefore the area of $ABCD$ is $\frac{63\sqrt{3}}{4} - 4\sqrt{3} = \frac{47\sqrt{3}}{4}$.
20. (B) Setting $x = 2$, and then $x = 1001$, we have $f(2) + 2f(1001) = 6$ and $f(1001) + 2f(2) = 3003$. Subtracting the first equation from twice the second equation we obtain $3f(2) = 6000$, so $f(2) = 2000$. Note that $f(x) = \frac{4004}{x} - x$ is such a function.
21. (C) Let $x = \log_c a$ and $y = \log_c b$. Then we are given

$$2\left(\frac{1}{x} + \frac{1}{y}\right) = \frac{9}{x+y}.$$

This is equivalent to $2(x+y)^2 = 9xy$ or

$$2x^2 - 5xy + 2y^2 = (2x - y)(x - 2y) = 0.$$

Therefore, $2\log_c a = \log_c b$ or $\log_c a = 2\log_c b$, and so $\log_a b = \frac{1}{2}$ or 2. The larger value is 2.

22. (D) If there are c ($c \geq 0$) correct answers and u ($u \geq 0$) unanswered questions and $c + u \leq 25$, then the score is $6c + 2.5u$. If c is sufficiently large and u is sufficiently small, the same score will be obtained with $c - 5$ correct answers and $u + 12$ unanswered questions (this requires $c + u \leq 18$), and also with $c - 10$ correct answers and $u + 24$ unanswered questions. Note that in the latter case we must have $c \geq 10$ and $c + u \leq 11$. Therefore, for there to be three ways to obtain the score $6c + 2.5u$ we can only have $c = 10$ and $u = 0$, or $c = 10$ and $u = 1$, or $c = 11$ and $u = 0$. The three such scores are 60, 62.5, and 66, and their sum is 188.5
23. (A) Let $z = w - 14i$. Then the equation becomes

$$(w - 14i)(w - 13i)(w - 11i) = 2002i.$$

This simplifies to

$$w^3 - 38iw^2 - 479w = 0.$$

The zeros of this cubic are 0 and $19i \pm \sqrt{118}$. Therefore the zeros of the original equation are $-14i$ and $5i \pm \sqrt{118}$, and the zero satisfying the given conditions is $\sqrt{118} + 5i$, so $a = \sqrt{118}$.

24. (B) Let a be the length of $ABCD$'s edges. Denote x, y, z the distances from E to the faces DAB, DBC, DCA respectively, and by u, v, w the distances from E to edges AB, BC, CA , respectively. Then

$$\begin{aligned} V_{ABCD} &= V_{EDAB} + V_{EDBC} + V_{EDCA} \\ &= \frac{1}{3}(K_{DAB} \cdot x + K_{DBC} \cdot y + K_{DCA} \cdot z) \end{aligned}$$

We have $V_{ABCD} = \frac{1}{3}K_{ABC} \cdot h$, where h is the altitude of tetrahedron $ABCD$. Then h is a leg in a right triangle whose other leg is $\frac{1}{3}a\frac{\sqrt{3}}{2} = \frac{a\sqrt{3}}{6}$ and whose hypotenuse is $\frac{a\sqrt{3}}{2}$. Then $h = \sqrt{\frac{3a^2}{4} - \frac{3a^2}{36}} = \frac{a\sqrt{6}}{3}$. Since ABC, DAB, DBC , and DCA are all equilateral triangles, $K_{ABC} = K_{DAB} = K_{DBC} = K_{DCA}$, so $s = x + y + z = h = \frac{a\sqrt{6}}{3}$. Similarly, in triangle ABC , $K_{ABC} = K_{EAB} + K_{EBC} + K_{ECA} = \frac{1}{2}(AB \cdot u + BC \cdot v + CA \cdot w)$. So $\frac{a^2\sqrt{3}}{4} = \frac{a}{2}(u + v + w)$. Hence, $S = u + v + w = \frac{a\sqrt{3}}{2}$ and $\frac{s}{S} = \frac{\frac{a\sqrt{6}}{3}}{\frac{a\sqrt{3}}{2}} = \frac{2\sqrt{2}}{3}$.

25. (C) Squaring each equation and adding gives

$$(\sin^2 a + \cos^2 a) + (\sin^2 b + \cos^2 b) + 2(\cos a \cos b + \sin a \sin b) = 2$$

or

$$\cos(a - b) = 0.$$

Multiplying the two equations together gives

$$(\sin a \cos b + \sin b \cos a) + (\sin a \cos a + \sin b \cos b) = \frac{\sqrt{3}}{2}$$

or

$$\sin(a + b) + \sin(a + b) \cos(a - b) = \frac{\sqrt{3}}{2}.$$

Substituting $\cos(a - b) = 0$, we have $\sin(a + b) = \frac{\sqrt{3}}{2}$.